

Distributed Nonsmooth Robust Resource Allocation with Cardinality Constrained Uncertainty

Yue Wei¹, Shuxin Ding², Hao Fang¹, Xianlin Zeng¹, Qingkai Yang¹, Bin Xin¹

1. School of Automation, Beijing Institute of Technology, Beijing 100081, P. R. China
E-mail: yuewei@bit.edu.cn; fangh@bit.edu.cn; xianlin.zeng@bit.edu.cn; qingkai.yang@bit.edu.cn; brucebin@bit.edu.cn

2. Signal and Communication Research Institute, China Academy of Railway Sciences, Beijing 100081, P. R. China
E-mail: jacketindd@gmail.com

Abstract: A distributed nonsmooth robust resource allocation problem with cardinality constrained uncertainty is investigated in this paper. The global objective is consisted of local objectives, which are convex but nonsmooth. Each agent is constrained in its private convex set and has only the information of its corresponding local objective. The resource allocation condition is subject to the cardinality constrained uncertainty sets. By employing the duality theory of convex optimization, a dual problem of the robust resource allocation problem is presented. For solving this dual problem, a distributed primal-dual projected algorithm is proposed. Theoretically, the convergence analysis by using stability theory of differential inclusions is conducted. It shows that the algorithm can steer the multi-agent system to satisfy resource allocation condition at the optimal solution. In the end, a nontrivial simulation is shown and the results demonstrate the efficiency of the proposed algorithm.

Key Words: Distributed Optimization, Robust Resource Allocation, Cardinality Constrained Uncertainty

1 Introduction

In recent years, the distributed optimization problem is widely studied as a hot topic in the areas of machine learning [1] and multi-agent system coordination [2]-[3]. In this problem, the objective is the sum of local objectives. Each agent can only obtain the knowledge of its private local objective. Many results of distributed optimization are focusing on steering the system to achieve consensus at the optimal solution [4]-[5]. On the other hand, the research of distributed globally constrained optimization has also gained a great of attention [6]-[7], especially the distributed resource allocation problem. In order to solve the distributed resource allocation problem, a distributed gradient-based algorithm was proposed while the initialization of states is required [8]. After this work, the initialization-free distributed algorithms for distributed resource allocation have been investigated in [9]-[10].

While most of the existing works about distributed resource allocation have the assumption that the resource allocation condition is deterministic. This assumption may not applied for the distributed resource allocation problems applying in the real world. In order to solve these problems, robustness of the distributed resource allocation should be stressed. Robust optimization deals with uncertainty described by uncertain-but-bounded parameters [11]. Typically, there are several kinds of uncertain parameters (eg., box/interval uncertainty, ellipsoidal uncertainty, polyhedral uncertainty, cardinality constrained uncertainty, etc.) [12]. Zeng et al. [6] proposed a distributed algorithm for robust resource allocation with polyhedral uncertain parameters. However, only considering polyhedral uncertain parameters may lead the problem too much conservative [13]. Cardinality constrained uncertainty provides a budget of uncertainty in terms of cardinality constraints which decrease the conservatism by combining interval and polyhedral uncertainty.

Besides, many real-world robust optimization problems are related with cardinality constrained uncertainty [14]. Therefore, the robust optimization problem with cardinality constrained uncertainty needs to be analysed.

Nonsmooth optimization problem is increasingly popular due to its important role in a lot of signal processing, statistical inference and machine learning problems. In the compressed sensing problem, the sparsity-promoting regulator has the form of l_1 -norm. In optimization problems with per-agent constraints, the indicator function of the constraint set of agent i is nonsmooth. In the geometric median problem, the objective is the mean of a sum of l_2 -norm functions.

In this paper, a distributed robust nonsmooth resource allocation problem with cardinality constrained uncertainty has been researched. The contributions of this paper are summed up as three parts:

- 1) The robust resource allocation problem we investigate here is with cardinality constrained uncertain parameters, which decrease the conservatism of the problem using polyhedral uncertain parameters.
- 2) We propose a distributed primal-dual projected algorithm with considering the duality theory of convex optimization.
- 3) The proof of the convergence of this algorithm has been given by employing the theory of nonsmooth analysis and differential inclusion.

This paper is organized as follows. Section 2 introduces the necessary preliminary concepts of graph theory, projection operator and differential inclusion. Section 3 shows the robust nonsmooth resource allocation problem with cardinality constrained uncertainty. Section 4 proposes a distributed projected primal-dual algorithm. Section 5 provides the convergence proof of the algorithm. Section 6 gives a numerical example to show the effectiveness of our proposed algorithm. Finally, we present conclusions in Section 7.

2 Preliminary

In this section, we introduce relevant notations, concepts on graph theory, projection operators and differential inclu-

This work is supported by Projects of Major International (Regional) Joint Research Program NSFC (Grant no. 61720106011), NSFC (Grant no. 61621063, 61573062, 61673058, 61603378, 61603094).

sions.

2.1 Graph Theory

A weighted undirected graph \mathcal{G} is denoted by $\mathcal{G}(\mathcal{V}, \mathcal{E}, \mathcal{A})$, where $\mathcal{V} = \{1, \dots, n\}$ is a set of nodes, $\mathcal{E} = \{(i, k) : i, k \in \mathcal{V}; i \neq k\} \subset \mathcal{V} \times \mathcal{V}$ is a set of edges, and $\mathcal{A} = [\alpha_{i,k}] \in \mathbb{R}^{n \times n}$ is a weighted adjacency matrix such that $\alpha_{i,k} = \alpha_{k,i} > 0$ if $(k, i) \in \mathcal{E}$ and $\alpha_{i,k} = 0$ otherwise, where $\mathbb{R}^{n \times n}$ denotes the set of n -by- n real matrices. $k \in \mathcal{N}_i$ denotes agent k is a neighbour of agent i . The Laplacian matrix is $L_n = D - \mathcal{A}$, where $D \in \mathbb{R}^{n \times n}$ is diagonal with $D_{i,i} = \sum_{k=1}^n \alpha_{i,k}$, $i \in \{1, \dots, n\}$. Specifically, if the weighted graph \mathcal{G} is undirected and connected, then $L_n = L_n^T \geq 0$.

2.2 Projection Operator

Define a projection operator as $P_\Omega(u) = \arg \min_{v \in \Omega} \|u - v\|$, where $\Omega \subset \mathbb{R}^n$ is closed and convex, \mathbb{R}^n denotes the set of n -dimensional real column vectors.

Lemma 2.1. [15] Let $\Omega \subset \mathbb{R}^n$ be closed and convex, and define $V : \mathbb{R}^n \rightarrow \mathbb{R}$ as $V(x) = \frac{1}{2}(\|x - P_\Omega(y)\|^2 - \|x - P_\Omega(x)\|^2)$ where $y \in \mathbb{R}^n$. Then $V(x) \geq \frac{1}{2}\|P_\Omega(x) - P_\Omega(y)\|^2$, $V(x)$ is differentiable and convex with respect to x , and $\nabla V(x) = P_\Omega(x) - P_\Omega(y)$.

Lemma 2.2. If $\Omega \subset \mathbb{R}^n$ is closed and convex, then $(P_\Omega(x) - P_\Omega(y))^T(x - y) \geq \|P_\Omega(x) - P_\Omega(y)\|^2$ for all $x, y \in \mathbb{R}^n$.

2.3 Differential Inclusion

Consider a nonsmooth system

$$\dot{x} \in \mathcal{F}(x(t)), x(0) = x_0, t \geq 0 \quad (1)$$

where $\mathcal{F} : \mathbb{R}^q \rightarrow \mathcal{B}(\mathbb{R}^q)$, $\mathcal{B}(\mathbb{R}^q)$ is the collection of subsets of \mathbb{R}^q . A set M is said to be weakly invariant (strongly invariant) with respect to (1) if for any $x_0 \in M$, M contains a maximal solution (all maximal solutions) of (1). An equilibrium point of (1) is a point $x^* \in \mathbb{R}^q$ such that $\mathbf{0}_q \in \mathcal{F}(x^*)$.

Let $V : \mathbb{R}^q \rightarrow \mathbb{R}$ be a locally Lipschitz continuous function and ∂V be the Clarke generalized gradient [16] of $V(x)$ at x . The set-valued Lie derivative [16] $\mathcal{L}_{\mathcal{F}}V : \mathbb{R}^q \rightarrow \mathcal{B}(\mathbb{R})$ of V with respect to (1) is defined as $\mathcal{L}_{\mathcal{F}}V(x) = \{a \in \mathbb{R} : \text{there exists } v \in \mathcal{F} \text{ such that } p^T v = a \text{ for all } p \in \partial V(x)\}$. In the case when $\mathcal{L}_{\mathcal{F}}V(x)$ is nonempty, we use $\max \mathcal{L}_{\mathcal{F}}V(x)$ to denote the largest element of $\mathcal{L}_{\mathcal{F}}V(x)$.

Lemma 2.3. [17] For the differential inclusion (1), we assume that \mathcal{F} is upper semicontinuous and locally bounded, and $\mathcal{F}(x)$ takes nonempty, compact, and convex values. Let $V : \mathbb{R}_q \rightarrow \mathbb{R}$ be a locally Lipschitz and regular function, $S \subset \mathbb{R}_q$ be compact and strongly invariant for (1), $\phi(\cdot)$ be a solution of (1), $\mathcal{R} = \{x \in \mathbb{R}_q : 0 \in \mathcal{L}_{\mathcal{F}}(x)\}$, and \mathcal{M} be the largest weakly invariant subset of $\bar{\mathcal{R}} \cap S$, where $\bar{\mathcal{R}}$ is the closure of \mathcal{R} . If $\max \mathcal{L}_{\mathcal{F}}V(x) \leq 0$ for all $x \in S$, then $\text{dist}(\phi(t), \mathcal{M}) \rightarrow 0$ as $t \rightarrow \infty$.

3 Problem Formulation

In this section, the distribute nonsmooth robust resource allocation optimization problem is formulated. Consider the following distributed nonsmooth uncertain resource allocation problem

$$\min_{x_i \in \Omega_i} f(x) = \sum_{i=1}^n f_i(x_i) \quad (2)$$

$$\text{s.t. } \sum_{i=1}^n \bar{a}_{ij}^l x_i^l \leq b_j^l, \quad \forall \bar{a}_{ij}^l \in \mathcal{U}_{ij}^l$$

where $j \in \{1, \dots, m\}$, $l \in \{1, \dots, q\}$, and with cardinality constrained uncertainty sets

$$\begin{aligned} \mathcal{U}_{ij}^l = & \{\bar{a}_{ij}^l | \bar{a}_{ij}^l \in [a_{ij}^l - \hat{a}_{ij}^l, a_{ij}^l + \hat{a}_{ij}^l], \\ & \sum_{i,l} \left| \frac{\bar{a}_{ij}^l - a_{ij}^l}{\hat{a}_{ij}^l} \right| \leq \gamma_j, \forall i, j, l\} \end{aligned} \quad (3)$$

For agent $i \in \{1, \dots, n\}$, $x_i \in \mathbb{R}^q$, Ω_i is the local constraint set, and $f_i(x_i)$ is the local objective which is continuous but not necessary smooth. $\bar{a}_{ij}^l \in \mathbb{R}$ is assumed to take arbitrary values in the uncertainty set \mathcal{U}_{ij}^l , $b_j = [b_j^1, \dots, b_j^q]^T \in \mathbb{R}^q$, and γ_j denotes the budget of uncertainty.

Then the corresponding robust optimization problem of the problem (2) is shown as

$$\begin{aligned} \min_{x_i \in \Omega_i} f(x) = & \sum_{i=1}^n f_i(x_i) \\ \text{s.t. } & \sum_{i=1}^n a_{ij}^l x_i^l + \max_{S_j^l \in \mathcal{J}_j^l: |S_j^l| = \gamma_j} \sum_{i \in S_j^l} \hat{a}_{ij}^l x_i^l \leq b_j^l, \\ & j \in \{1, \dots, m\}, l \in \{1, \dots, q\} \end{aligned} \quad (4)$$

For the l -th dimensional elements of each agent's states with the j -th resource allocation condition, S_j^l is a possible set of the chosen agents where the size of S_j^l is γ_j , and \mathcal{J}_j^l is the set of all possible S_j^l .

According to the duality of convex optimization [18], the problem (4) can be transferred to the corresponding dual problem as

$$\begin{aligned} \min_{x_i \in \Omega_i} f(x) = & \sum_{i=1}^n f_i(x_i) \\ \text{s.t. } & \sum_{i=1}^n [A_{ij} x_i + \frac{1}{n} \gamma_j z_{ij} + w_{ij}] \leq \sum_{i=1}^n b_{ij}, \\ & \hat{A}_{ij} x_i \leq z_{ij} + w_{ij}, L_{mnq} Z = \mathbf{0}_{mnq}, \\ & z_{ij} \geq \mathbf{0}_q, w_{ij} \geq \mathbf{0}_q, \\ & i \in \{1, \dots, n\}, j \in \{1, \dots, m\} \end{aligned} \quad (5)$$

where $A_{ij} = \text{diag}\{a_{ij}^1, \dots, a_{ij}^q\} \in \mathbb{R}^{q \times q}$, $\hat{A}_{ij} = \text{diag}\{\hat{a}_{ij}^1, \dots, \hat{a}_{ij}^q\} \in \mathbb{R}^{q \times q}$, $z_{ij} \in \mathbb{R}^q$, $z_i = [(z_{i1})^T, \dots, (z_{im})^T]^T$, $Z = [(z_1)^T, \dots, (z_n)^T]^T$, $w_{ij} \in \mathbb{R}^q$, $\sum_{i=1}^n b_{ij} = b_j$, $L_{mnq} = L_n \otimes I_{mq}$, where $A \otimes B$ denotes the Kronecker product of matrices A and B . $\mathbf{0}_n$ is the $n \times 1$ vector with all elements of 0.

The assumptions below are made for the wellposedness of the problem (5) in this section.

Assumption 3.1. 1) The weighted graph \mathcal{G} is connected and undirected.

2) For $i \in \{1, \dots, n\}$, f_i is strictly convex on an open set containing Ω_i , and $\Omega_i \subset \mathbb{R}^q$ is closed and convex.

3) (Slater's constraint condition) There exist $x_i \in \Omega_i$, $z_{ij} \in \mathbb{R}_+^q$ and $w_{ij} \in \mathbb{R}_+^q$ satisfying the constraint for

$i \in \{1, \dots, n\}$ and $j \in \{1, \dots, m\}$, where $\bar{\mathbb{R}}_+^q$ denotes the set of nonnegative q -dimensional real column vectors.

The Lagrangian of dual problem (5) is described as

$$\begin{aligned} & L(x, Z, W, \Lambda^1, \Lambda^2, U) \\ &= \sum_{i=1}^n [f^i(x_i) + \sum_{j=1}^m [(\lambda_{ij}^1)^T (A_{ij}x_i + \frac{1}{n}\gamma_j z_{ij} + w_{ij} - b_{ij}) \\ & \quad + (\lambda_{ij}^2)^T (\hat{A}_{ij}x_i - z_{ij} - w_{ij})]] + \mu^T L_{mnq}Z \end{aligned} \quad (6)$$

where $w_i = [(w_{i1})^T, \dots, (w_{im})^T]^T$, $W = [(w_1)^T, \dots, (w_n)^T]^T$, $\lambda_i^g = [(\lambda_{i1}^g)^T, \dots, (\lambda_{im}^g)^T]^T$, $\Lambda^g = [(\lambda_1^g)^T, \dots, (\lambda_n^g)^T]^T$, $\mu_i = [(\mu_{i1})^T, \dots, (\mu_{im})^T]^T$, $U = [\mu_1^T, \dots, \mu_n^T]^T$, $g \in \{1, 2\}$.

Then according to problem (5), the following lemma is arrived by the Karush-Kuhn-Tucker (KKT) condition of convex optimization problems.

Lemma 3.1. Under the Assumptions 3.1, a feasible point $x^* \in \mathbb{R}^{nq}$ is a minimizer to Problem (5) if and only if there exist $x_i^* \in \Omega_i \in \mathbb{R}^q$, $\lambda_{ij}^{1*} \in \mathbb{R}^q$, $\lambda_{ij}^{2*} \in \mathbb{R}^q$, $\mu_{ij}^* \in \mathbb{R}^q$, $w_{ij}^* \in \mathbb{R}^q$ and $z_{ij}^* \in \mathbb{R}^q$ such that for $i \in \{1, \dots, n\}$, $j \in \{1, \dots, m\}$

$$-\partial f_i(x_i^*) - \sum_{j=1}^m A_{ij}\lambda_{ij}^{1*} - \sum_{j=1}^m \hat{A}_{ij}\lambda_{ij}^{2*} \in \mathcal{N}_{\Omega_i}(x_i^*), \quad (7a)$$

$$-\frac{1}{n}\gamma_j\lambda_{ij}^{1*} + \lambda_{ij}^{2*} - \sum_{k \in \mathcal{N}_i} (\mu_{ij} - \mu_{kj}) \in \mathcal{N}_{\bar{\mathbb{R}}_+^q}(z_{ij}^*), \quad (7b)$$

$$-\lambda_{ij}^{1*} + \lambda_{ij}^{2*} \in \mathcal{N}_{\bar{\mathbb{R}}_+^q}(w_{ij}^*), \quad (7c)$$

$$\sum_{i=1}^n [A_{ij}x_i^* + \frac{1}{n}\gamma_j z_{ij}^* + w_{ij}^* - b_{ij}] \leq \mathbf{0}_q, \quad (7d)$$

$$\sum_{i=1}^n [\hat{A}_{ij}x_i^* - z_{ij}^* - w_{ij}^*] \leq \mathbf{0}_q, \quad (7e)$$

$$L_{mnq}Z^* = \mathbf{0}_{mnq}, \quad (7f)$$

$$(\lambda_{ij}^{1*})^T [A_{ij}x_i^* + \frac{1}{n}\gamma_j z_{ij}^* + w_{ij}^* - b_{ij}] = 0, \quad (7g)$$

$$(\lambda_{ij}^{2*})^T [\hat{A}_{ij}x_i^* - z_{ij}^* - w_{ij}^*] = 0. \quad (7h)$$

where $\mathcal{N}_{\Omega_i}(x_i^*)$ is the the normal cone of Ω_i at x_i^* .

The proof of Lemma 3.1 is omitted since it is a trivial extension of the proof for Theorem 3.34 in [19].

4 Algorithm Design

In this section, we propose a distributed algorithm for this problem (5). The algorithm is detailed as below:

$$\begin{cases} \dot{\hat{x}}_i \in -\bar{x}_i + x_i - \partial f_i(x_i) - \sum_{j=1}^m A_{ij}\lambda_{ij}^1 - \sum_{j=1}^m \hat{A}_{ij}\lambda_{ij}^2 \\ \dot{\hat{z}}_{ij} = -\bar{z}_{ij} + z_{ij} - \frac{1}{n}\gamma_j\lambda_{ij}^1 + \lambda_{ij}^2 - \sum_{k \in \mathcal{N}_i} \alpha_k(\mu_{ij} - \mu_{kj}) \\ \dot{\hat{w}}_{ij} = -\bar{w}_{ij} + w_{ij} - \lambda_{ij}^1 + \lambda_{ij}^2 \\ \dot{\hat{\mu}}_{ij} = \sum_{k \in \mathcal{N}_i} \alpha_{ik}(z_{ij} - z_{kj}) \\ \dot{\hat{\lambda}}_{ij}^1 = -\bar{\lambda}_{ij}^1 + \lambda_{ij}^1 + [A_{ij}x_i + \frac{1}{n}\gamma_j z_{ij} + w_{ij} - b_{ij}] \\ \quad + \sum_{k \in \mathcal{N}_i} \alpha_{ik}(y_{ij}^1 - y_{kj}^1) - \sum_{k \in \mathcal{N}_i} \alpha_{ik}(\lambda_{ij}^1 - \lambda_{kj}^1) \\ \dot{\hat{\lambda}}_{ij}^2 = -\bar{\lambda}_{ij}^2 + \lambda_{ij}^2 + [\hat{A}_{ij}x_i - z_{ij} - w_{ij}] \\ \quad + \sum_{k \in \mathcal{N}_i} \alpha_{ik}(y_{ij}^2 - y_{kj}^2) - \sum_{k \in \mathcal{N}_i} \alpha_{ik}(\lambda_{ij}^2 - \lambda_{kj}^2) \\ \dot{y}_{ij}^1 = -\sum_{k \in \mathcal{N}_i} \alpha_{ik}(\lambda_{ij}^1 - \lambda_{kj}^1) \\ \dot{y}_{ij}^2 = -\sum_{k \in \mathcal{N}_i} \alpha_{ik}(\lambda_{ij}^2 - \lambda_{kj}^2) \\ x_i = P_{\Omega_i}[\bar{x}_i], z_{ij} = P_{\bar{\mathbb{R}}_+^q}[\bar{z}_{ij}], w_{ij} = P_{\bar{\mathbb{R}}_+^q}[\bar{w}_{ij}], \\ \lambda_{ij}^1 = P_{\bar{\mathbb{R}}_+^q}[\bar{\lambda}_{ij}^1], \lambda_{ij}^2 = P_{\bar{\mathbb{R}}_+^q}[\bar{\lambda}_{ij}^2] \end{cases} \quad (8)$$

where $t \geq 0$.

The algorithm (8) can be also written as a compact form as

$$\begin{aligned} \dot{\Phi} &\in \mathcal{F}(\Phi), x = P_{\Omega}[\bar{x}], Z = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{Z}], \\ W &= P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{W}], \Lambda^1 = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{\Lambda}^1], \Lambda^2 = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{\Lambda}^2] \end{aligned} \quad (9)$$

where $\Phi = [\bar{x}^T, \bar{Z}^T, \bar{W}^T, U^T, (\bar{\Lambda}^1)^T, (\bar{\Lambda}^2)^T, (Y^1)^T, (Y^2)^T]^T$, $P_{\Omega}[\bar{x}] = [(P_{\Omega_1}[\bar{x}_1])^T, \dots, (P_{\Omega_n}[\bar{x}_n])^T]^T$, $\bar{z}_i = [(\bar{z}_{i1})^T, \dots, (\bar{z}_{im})^T]^T$, $\bar{Z} = [(\bar{z}_1)^T, \dots, (\bar{z}_n)^T]^T$, $\bar{w}_i = [(\bar{w}_{i1})^T, \dots, (\bar{w}_{im})^T]^T$, $\bar{W} = [(\bar{w}_1)^T, \dots, (\bar{w}_n)^T]^T$, $\bar{\lambda}_i^g = [(\bar{\lambda}_{i1}^g)^T, \dots, (\bar{\lambda}_{im}^g)^T]^T$, $\bar{\Lambda}^g = [(\bar{\lambda}_1^g)^T, \dots, (\bar{\lambda}_n^g)^T]^T$, $y_i^g = [(y_{i1}^g)^T, \dots, (y_{im}^g)^T]^T$, $Y^g = [(y_1^g)^T, \dots, (y_n^g)^T]^T$, $g \in \{1, 2\}$.

In (9), $\mathcal{F}(\phi)$ is defined as

$$\begin{aligned} \mathcal{F}(\phi) &= \{[p_{\bar{x}}^T, p_{\bar{Z}}^T, p_{\bar{W}}^T, p_U^T, p_{\bar{\Lambda}^1}^T, p_{\bar{\Lambda}^2}^T, p_{Y^1}^T, p_{Y^2}^T]^T \\ &\in \mathbb{R}^{nq} \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq} \\ &\quad \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq}\} \end{aligned} \quad (10)$$

with

$$\begin{cases} p_{\bar{x}} = -\bar{x} + x - f_x - EA^*\Lambda^1 - E\hat{A}^*\Lambda^2 \\ p_{\bar{Z}} = -\bar{Z} + Z - \frac{1}{n}\Gamma\Lambda^1 + \Lambda^2 - L_{mnq}U \\ p_{\bar{W}} = -\bar{W} + W - \Lambda^1 + \Lambda^2, p_U = L_{mnq}Z \\ p_{\bar{\Lambda}^1} = -\bar{\Lambda}^1 + \Lambda^1 + [A^*E^T x + \frac{1}{n}\Gamma Z + W - B] \\ \quad + L_{mnq}Y^1 - L_{mnq}\Lambda^1 \\ p_{\bar{\Lambda}^2} = -\bar{\Lambda}^2 + \Lambda^2 + [\hat{A}^*E^T x - Z - W] \\ \quad + L_{mnq}Y^2 - L_{mnq}\Lambda^2 \\ p_{Y^1} = -L_{mnq}\Lambda^1, p_{Y^2} = -L_{mnq}\Lambda^2 \end{cases} \quad (11)$$

where $E = I_n \otimes (\mathbf{1}_m \otimes I_q) \in \mathbb{R}^{nq \times mnq}$, I_n is the n -dimensional identity matrix, $\mathbf{1}_m$ denotes the $n \times 1$ vector with all elements of 1. $A_i = \text{diag}\{A_{i1}, \dots, A_{im}\} \in \mathbb{R}^{mq \times mq}$, $A^* = \text{diag}\{A_1, \dots, A_n\} \in \mathbb{R}^{mnq \times mnq}$, $\hat{A}_i = \text{diag}\{\hat{A}_{i1}, \dots, \hat{A}_{im}\} \in \mathbb{R}^{mq \times mq}$, $\hat{A}^* = \text{diag}\{\hat{A}_1, \dots, \hat{A}_n\} \in \mathbb{R}^{mnq \times mnq}$, $\Gamma = I_n \otimes \text{diag}\{\gamma_1 I_q, \dots, \gamma_m I_q\} \in \mathbb{R}^{mnq \times mnq}$, $L_{mnq} = L_n \otimes I_{mq}$, $b_i = [(b_{i1})^T, \dots, (b_{im})^T]^T$, $B = [(b_1)^T, \dots, (b_n)^T]^T$, $f_x \in \partial f(x)$.

Then the equilibrium of algorithm (9) is

$$\mathbf{0}_{nq} = -\bar{x}^* + x^* - f_{x^*} - EA^*\Lambda^{1*} - E\hat{A}^*\Lambda^{2*} \quad (12a)$$

$$\mathbf{0}_{mnq} = -\bar{Z}^* + Z^* - \frac{1}{n}\Gamma\Lambda^{1*} + \Lambda^{2*} - L_{mnq}U^* \quad (12b)$$

$$\mathbf{0}_{mnq} = -\bar{W}^* + W^* - \Lambda^{1*} + \Lambda^{2*} \quad (12c)$$

$$\mathbf{0}_{mnq} = L_{mnq}Z^* \quad (12d)$$

$$\mathbf{0}_{mnq} = -\bar{\Lambda}^{1*} + \Lambda^{1*} + [A^*E^T x^* + \frac{1}{n}\Gamma Z^* + W^* - B] + L_{mnq}Y^{1*} \quad (12e)$$

$$\mathbf{0}_{mnq} = -\bar{\Lambda}^{2*} + \Lambda^{2*} + [\hat{A}^*E^T x^* - Z^* - W^*] + L_{mnq}Y^{2*} \quad (12f)$$

$$\mathbf{0}_{mnq} = L_{mnq}\Lambda^{1*}, \mathbf{0}_{mnq} = L_{mnq}\Lambda^{2*} \quad (12g)$$

$$x^* = P_{\Omega}[\bar{x}^*], Z^* = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{Z}^*], W^* = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{W}^*] \quad (12h)$$

$$\Lambda^{1*} = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{\Lambda}^{1*}], \Lambda^{2*} = P_{\bar{\mathbb{R}}_+^{mnq}}[\bar{\Lambda}^{2*}] \quad (12i)$$

Here we give the Lemma 4.1 to link the equilibrium of algorithm with the solution of problem (4).

Lemma 4.1. Consider Problem (4) and Assumption 3.1 holds. If $\phi^* \in \mathbb{R}^{(7m+1)nq}$ is an equilibrium of (8), then $x^* = P_\Omega[\bar{x}^*]$ is a solution to Problem (4).

Proof. Suppose $\phi^* \in \mathbb{R}^{(7m+1)nq}$ is an equilibrium of (8). When considering (12a), (12b) and (12c), there exists $f_{x^*} \in \partial f(x^*)$ such that $\bar{x}^* = x^* - f_{x^*} - EA^*\Lambda^{1*} - E\hat{A}^*\Lambda^{2*}$, $\bar{Z}^* = Z^* - \frac{1}{n}\Gamma\Lambda^{1*} + \Lambda^{2*} - L_{mnq}U^*$, $\bar{W}^* = W^* - \Lambda^{1*} + \Lambda^{2*}$. Since $x^* = P_\Omega[\bar{x}^*]$, $Z^* = P_{\mathbb{R}_+^{mnq}}[\bar{Z}^*]$, $W^* = P_{\mathbb{R}_+^{mnq}}[\bar{W}^*]$, it follows that (7a), (7b) and (7c) holds.

According to (12e) and (12f), one can have that

$$Q_j(-\bar{\Lambda}^{1*} + \Lambda^{1*} + [A^*E^T x^* + \frac{1}{n}\Gamma Z^* + W^* - B] + L_{mnq}Y^{1*}) = -\sum_{i=1}^n(\bar{\lambda}_{ij}^{1*} - \lambda_{ij}^{1*}) + \sum_{i=1}^n H_{ij}^{1*} = \mathbf{0}_q \quad (13)$$

$$Q_j(-\bar{\Lambda}^{2*} + \Lambda^{2*} + [\hat{A}^*E^T x^* - Z^* - W^*] + L_{mnq}Y^{2*}) = -\sum_{i=1}^n(\bar{\lambda}_{ij}^{2*} - \lambda_{ij}^{2*}) + \sum_{i=1}^n H_{ij}^{2*} = \mathbf{0}_q \quad (14)$$

where $Q_j = \mathbf{1}_n^T \otimes (I_m^j \otimes I_q) \in \mathbb{R}^{q \times mnq}$, I_m^j denotes the j -th row of I_m . $Q_j L_{mnq} Y^{1*} = \mathbf{0}_q$, $Q_j L_{mnq} Y^{2*} = \mathbf{0}_q$, $H_{i,j}^1 = A_{ij}x_i + \frac{1}{n}\gamma_j z_{ij} + w_{ij} - b_{ij}$, $H_{i,j}^2 = \hat{A}_{ij}x_i - z_{ij} - w_{ij}$. Since $\lambda_{ij}^1 = P_{\mathbb{R}_+^{mnq}}[\bar{\lambda}_{ij}^1] \geq \mathbf{0}_q$, $\lambda_{ij}^2 = P_{\mathbb{R}_+^{mnq}}[\bar{\lambda}_{ij}^2] \geq \mathbf{0}_q$, $\bar{\lambda}_{ij}^1 - \lambda_{ij}^1 \leq \mathbf{0}_q$ and $\bar{\lambda}_{ij}^2 - \lambda_{ij}^2 \leq \mathbf{0}_q$ for all $i \in \{1, \dots, n\}$. Hence (7d) and (7e) holds. (12d) equals to (7f), which means that (7f) holds.

It follows from (12g), (12i) and $\lambda_{ij}^1 = P_{\mathbb{R}_+^{mnq}}[\bar{\lambda}_{ij}^1] \geq \mathbf{0}_q$, $\lambda_{ij}^2 = P_{\mathbb{R}_+^{mnq}}[\bar{\lambda}_{ij}^2] \geq \mathbf{0}_q$ that there exist $\lambda_0^{1*} \in \mathbb{R}_+^{mq}$ and $\lambda_0^{2*} \in \mathbb{R}_+^{mq}$ such that $\Lambda^{1*} = \lambda_0^{1*} \otimes \mathbf{1}_n$ and $\Lambda^{2*} = \lambda_0^{2*} \otimes \mathbf{1}_n$. If $\lambda_0^{1*} = \mathbf{0}_{mq}$ and $\lambda_0^{2*} = \mathbf{0}_{mq}$, the (7g) and (7h) holds. If $\lambda_0^{1*} > \mathbf{0}_{mq}$ and $\lambda_0^{2*} > \mathbf{0}_{mq}$, it is clear that $\hat{\lambda}_{ij}^{1*} = \lambda_{ij}^{1*}$, $\hat{\lambda}_{ij}^{2*} = \lambda_{ij}^{2*}$, $\sum_{i=1}^n H_{ij}^{1*} = \mathbf{0}_q$ and $\sum_{i=1}^n H_{ij}^{2*} = \mathbf{0}_q$, which means that (7g) and (7h) also holds.

By Lemma 3.1, $(x^*, Z^*, W^*) \in \Omega \times \mathbb{R}^{mnq} \times \mathbb{R}^{mnq}$ is an optimal solution of Problem (5). Note that Problem (5) is the strong dual problem of Problem (4). Then the proof is accomplished. \square

5 Main Result

In this section, we give the convergence analysis of our algorithm (8). Define the Lyapunov candidate

$$V(\phi) = V_1(\bar{x}) + V_2(\bar{Z}) + V_3(\bar{W}) + V_4(U) + V_5(\bar{\Lambda}^1) + V_6(\bar{\Lambda}^2) + V_7(Y^1) + V_8(Y^2) \quad (15)$$

where

$$\begin{cases} V_1(\bar{x}) = \frac{1}{2}(\|\bar{x} - x^*\|^2 - \|\bar{x} - x\|^2) \\ V_2(\bar{Z}) = \frac{1}{2}(\|\bar{Z} - Z^*\|^2 - \|\bar{Z} - Z\|^2) \\ V_3(\bar{W}) = \frac{1}{2}(\|\bar{W} - W^*\|^2 - \|\bar{W} - W\|^2) \\ V_4(U) = \frac{1}{2}(\|U - U^*\|^2) \\ V_5(\bar{\Lambda}^1) = \frac{1}{2}(\|\bar{\Lambda}^1 - \Lambda^{1*}\|^2 - \|(\bar{\Lambda}^1 - \Lambda^{1*})\|^2) \\ V_6(\bar{\Lambda}^2) = \frac{1}{2}(\|\bar{\Lambda}^2 - \Lambda^{2*}\|^2 - \|(\bar{\Lambda}^2 - \Lambda^{2*})\|^2) \\ V_7(Y^1) = \frac{1}{2}(\|Y^1 - Y^{1*}\|^2) \\ V_8(Y^2) = \frac{1}{2}(\|Y^2 - Y^{2*}\|^2) \end{cases} \quad (16)$$

In the following lemma, we have analysed the set-valued derivative of $V(\phi)$ defined in (15) along the trajectories of Algorithm (8).

Lemma 5.1. Consider Algorithm (8) under Assumption 3.1 with $V(\phi)$ defined in (15). If $\beta \in \mathcal{L}_{\mathcal{F}}V(\phi)$, then there exist $f_x \in \partial f(x)$ and $f_{x^*} \in \partial f(x^*)$ with $x = P_\Omega[\bar{x}]$ and $x^* = P_\Omega[\bar{x}^*]$ such that

$$\beta \leq -(x - x^*)^T(f_x - f_{x^*}) - (\Lambda^1)^T L_{mnq} \Lambda^1 - (\Lambda^2)^T L_{mnq} \Lambda^2 \leq 0 \quad (17)$$

Proof. It follows from Lemma 2.1 that the gradients of $V(\phi)$ with respect to ϕ are

$$\begin{cases} \nabla_{\bar{x}} V(\phi) = x - x^*, \nabla_{\bar{Z}} V(\phi) = Z - Z^* \\ \nabla_{\bar{W}} V(\phi) = W - W^*, \nabla_U V(\phi) = U - U^* \\ \nabla_{\bar{\Lambda}^1} V(\phi) = \Lambda^1 - \Lambda^{1*}, \nabla_{\bar{\Lambda}^2} V(\phi) = \Lambda^2 - \Lambda^{2*} \\ \nabla_{Y^1} V(\phi) = Y^1 - Y^{1*}, \nabla_{Y^2} V(\phi) = Y^2 - Y^{2*} \end{cases} \quad (18)$$

The function $V(\phi)$ with the trajectories of (8) satisfies

$$\begin{aligned} \mathcal{L}_{\mathcal{F}}V(\phi) = & \{\beta \in \mathbb{R} : \beta = \nabla_{\bar{x}} V(\phi)^T p_{\bar{x}} + \nabla_{\bar{Z}} V(\phi)^T p_{\bar{Z}} \\ & + \nabla_{\bar{W}} V(\phi)^T p_{\bar{W}} + \nabla_U V(\phi)^T p_U \\ & + \nabla_{\bar{\Lambda}^1} V(\phi)^T p_{\bar{\Lambda}^1} + \nabla_{\bar{\Lambda}^2} V(\phi)^T p_{\bar{\Lambda}^2} \\ & + \nabla_{Y^1} V(\phi)^T p_{Y^1} + \nabla_{Y^2} V(\phi)^T p_{Y^2}\} \end{aligned} \quad (19)$$

Suppose $\beta \in \mathcal{L}_{\mathcal{F}}V(\phi)$. There exists $f_x \in \partial f(x)$ such that $\beta = \sum_{i=1}^8 \beta_i$, then

$$\begin{cases} \beta_1 = (x - x^*)^T(-\bar{x} + x - f_x - EA^*\Lambda^{1*} - E\hat{A}^*\Lambda^{2*}) \\ \beta_2 = (Z - Z^*)^T(-\bar{Z} + Z - \frac{1}{n}\Gamma\Lambda^{1*} + \Lambda^{2*} - L_{mnq}U) \\ \beta_3 = (W - W^*)^T(-\bar{W} + W - \Lambda^{1*} + \Lambda^{2*}) \\ \beta_4 = (U - U^*)^T L_{mnq} Z \\ \beta_5 = (\Lambda^1 - \Lambda^{1*})^T(-\bar{\Lambda}^1 + \Lambda^1 + [A^*E^T x + \frac{1}{n}\Gamma Z + W - B] + L_{mnq}Y^1 - L_{mnq}\Lambda^1) \\ \beta_6 = (\Lambda^2 - \Lambda^{2*})^T(-\bar{\Lambda}^2 + \Lambda^2 + [\hat{A}^*E^T x - Z - W] + L_{mnq}Y^2 - L_{mnq}\Lambda^2) \\ \beta_7 = -(Y^1 - Y^{1*})^T L_{mnq} \Lambda^1 \\ \beta_8 = -(Y^2 - Y^{2*})^T L_{mnq} \Lambda^2 \end{cases} \quad (20)$$

Since $\phi^* \in \mathbb{R}^{(7m+1)nq}$ is an equilibrium of (8), there exists $f_{x^*} \in \partial f(x^*)$ such that (12) holds.

From (12) and (20), one can have that

$$\begin{aligned} \beta = & -(x - x^*)^T(\bar{x} - \bar{x}^*) + \|x - x^*\|^2 \\ & - (Z - Z^*)^T(\bar{Z} - \bar{Z}^*) + \|Z - Z^*\|^2 \\ & - (W - W^*)^T(\bar{W} - \bar{W}^*) + \|W - W^*\|^2 \\ & - (\Lambda^1 - \Lambda^{1*})^T(\bar{\Lambda}^1 - \bar{\Lambda}^{1*}) + \|\Lambda^1 - \Lambda^{1*}\|^2 \\ & - (\Lambda^2 - \Lambda^{2*})^T(\bar{\Lambda}^2 - \bar{\Lambda}^{2*}) + \|\Lambda^2 - \Lambda^{2*}\|^2 \\ & - (\Lambda^1)^T L_{mnq} \Lambda^1 - (\Lambda^2)^T L_{mnq} \Lambda^2 \\ & - (x - x^*)^T(f_x - f_{x^*}) \end{aligned} \quad (21)$$

Since $L_{mnq} \geq 0$ and $(x - x^*)^T(f_x - f_{x^*}) \geq 0$ followed by the convexity of f , then according to Lemma 2.2, it follows from (21) that (17) is satisfied. \square

The following theorem proves the convergence of trajectory $x(t)$ with the proposed algorithm (8) to the optimal solutions.

Theorem 5.1. For Algorithm (8) with Assumption 3.1, we have that the results that

- (i) the trajectory $(x, Z, W, \Lambda^1, \Lambda^2, \phi)$ is bounded;
- (ii) $x(t)$ converges to the optimal solution to Problem (4).

Proof. i) Let $V(\phi)$ be as defined in (15). It follows from Lemma 5.1 that

$$\begin{aligned} & \max \mathcal{L}_{\mathcal{F}} V(\phi) \\ & \leq \max \{ -(x - x^*)^T (f_x - f_{x^*}) - (\Lambda^1)^T L_{mnq} \Lambda^1 \\ & \quad - (\Lambda^2)^T L_{mnq} \Lambda^2 : f_x \in \partial f(x), f_{x^*} \in \partial f(x^*), \quad (22) \\ & \quad x = P_{\Omega}[\bar{x}], \Lambda^1 = P_{\mathbb{R}_+^{mnq}}[\bar{\Lambda}^1], \Lambda^2 = P_{\mathbb{R}_+^{mnq}}[\bar{\Lambda}^2] \} \leq 0 \end{aligned}$$

Note that $V(\phi) \geq \frac{1}{2}(\|x - x^*\|^2 + \|Z - Z^*\|^2 + \|W - W^*\|^2 + \|U - U^*\|^2 + \|\Lambda^1 - \Lambda^{1*}\|^2 + \|\Lambda^2 - \Lambda^{2*}\|^2 + \|Y^1 - Y^{1*}\|^2 + \|Y^2 - Y^{2*}\|^2)$ according to Lemma 2.1. Hence that trajectory $(x(t), Z(t), W(t), U(t), \Lambda^1(t), \Lambda^2(t), Y^1(t), Y^2(t))$, $t \geq 0$ is bounded.

Because $\partial f(x)$ is compact for all $x \in \Omega$ and $(x(t), Z(t), W(t), U(t), \Lambda^1(t), \Lambda^2(t), Y^1(t), Y^2(t))$ is bounded for all $t \geq 0$, there exists $M = M(x, Z, W, \Lambda^1, \Lambda^2, \phi) > 0$ such that

$$\begin{cases} M \geq \|x(t) - f_{x(t)} - EA^* \Lambda^1(t) - E\hat{A}^* \Lambda^2(t)\| \\ M \geq \|Z(t) - \frac{1}{n} \Gamma \Lambda^1(t) + \Lambda^2(t) - L_{mnq} U(t)\| \\ M \geq \|W(t) - \Lambda^1(t) + \Lambda^2(t)\| \\ M \geq \|\Lambda^1(t) + [A^* E^T x(t) + \frac{1}{n} \Gamma Z(t) + W(t) \\ \quad - B] + L_{mnq} Y^1(t) - L_{mnq} \Lambda^1\| \\ M \geq \|\Lambda^2(t) + [\hat{A}^* E^T x(t) - Z(t) - W(t) \\ \quad + L_{mnq} Y^2(t) - L_{mnq} \Lambda^2(t)]\| \end{cases} \quad (23)$$

for all $f_{x(t)} \in \partial f(x(t))$ and all $t \geq 0$. Define $X : \mathcal{R}^{nq} \times \mathcal{R}^{mnq} \times \mathcal{R}^{mnq} \times \mathcal{R}^{mnq} \times \mathcal{R}^{mnq} \rightarrow \mathcal{R}$ by

$$X(\bar{x}, \bar{Z}, \bar{W}, \bar{\Lambda}^1, \bar{\Lambda}^2) = \frac{1}{2} (\|\bar{x}\|^2 + \|\bar{Z}\|^2 + \|\bar{W}\|^2 + \sum_{i=1}^2 \|\bar{\Lambda}^i\|^2) \quad (24)$$

The function $X(\bar{x}, \bar{Z}, \bar{W}, \bar{\Lambda}^1, \bar{\Lambda}^2)$ along the trajectories of (8) satisfies that

$$\begin{aligned} & \mathcal{L}_{\mathcal{F}} X(\bar{x}, \bar{Z}, \bar{W}, \bar{\Lambda}^1, \bar{\Lambda}^2) \\ & = \{x^T(-\bar{x} + x - f_x - EA^* \Lambda^1 - E\hat{A}^* \Lambda^2) \\ & \quad + \bar{Z}^T(-\bar{Z} + Z - \frac{1}{n} \Gamma \Lambda^1 + \Lambda^2 - L_{mnq} U) \\ & \quad + \bar{W}^T(-\bar{W} + W - \Lambda^1 + \Lambda^2) + (\bar{\Lambda}^1)^T(-\bar{\Lambda}^1 + \Lambda^1 \\ & \quad + [A^* E^T x + \frac{1}{n} \Gamma Z + W - B] + L_{mnq} Y^1 - L_{mnq} \Lambda^1) \\ & \quad + (\bar{\Lambda}^2)^T(-\bar{\Lambda}^2 + \Lambda^2 + [\hat{A}^* E^T x - Z - W] + L_{mnq} Y^2 \\ & \quad - L_{mnq} \Lambda^2) : f_x \in \partial f(x), x = P_{\Omega}[\bar{x}], Z = P_{\mathbb{R}_+^{mnq}}[\bar{Z}], \\ & \quad W = P_{\mathbb{R}_+^{mnq}}[\bar{W}], \Lambda^1 = P_{\mathbb{R}_+^{mnq}}[\bar{\Lambda}^1], \Lambda^2 = P_{\mathbb{R}_+^{mnq}}[\bar{\Lambda}^2] \} \end{aligned} \quad (25)$$

Note that

$$\begin{cases} -\|\bar{x}\|^2 + M\|\bar{x}\| \geq x^T(t)(-\bar{x}(t) + x(t) - f_{x(t)} \\ \quad - EA^* \Lambda^1(t) - E\hat{A}^* \Lambda^2(t)) \\ -\|\bar{Z}\|^2 + M\|\bar{Z}\| \geq \bar{Z}^T(t)(-\bar{Z}(t) + Z(t) - \frac{1}{n} \Gamma \Lambda^1(t) \\ \quad + \Lambda^2(t) - L_{mnq} U(t)) \\ -\|\bar{W}\|^2 + M\|\bar{W}\| \geq \bar{W}^T(t)(-\bar{W}(t) + W(t) \\ \quad - \Lambda^1(t) + \Lambda^2(t)) \\ -\|\bar{\Lambda}^1\|^2 + M\|\bar{\Lambda}^1\| \geq (\bar{\Lambda}^1)^T(t)(-\bar{\Lambda}^1(t) + \Lambda^1(t) \\ \quad + [A^* E^T x(t) + \frac{1}{n} \Gamma Z(t) + W(t) \\ \quad - B] + L_{mnq} Y^1(t) - L_{mnq} \Lambda^1(t)) \\ -\|\bar{\Lambda}^2\|^2 + M\|\bar{\Lambda}^2\| \geq (\bar{\Lambda}^2)^T(t)(-\bar{\Lambda}^2(t) + \Lambda^2(t) \\ \quad + [\hat{A}^* E^T x(t) - Z(t) - W(t)] \\ \quad + L_{mnq} Y^2(t) - L_{mnq} \Lambda^2(t)) \end{cases} \quad (26)$$

Hence,

$$\begin{aligned} & \max \mathcal{L}_{\mathcal{F}} X(\bar{x}(t), \bar{Z}(t), \bar{W}(t), \bar{\Lambda}^1(t), \bar{\Lambda}^2(t)) \\ & \leq -2X(\bar{x}(t), \bar{Z}(t), \bar{W}(t), \bar{\Lambda}^1(t), \bar{\Lambda}^2(t)) \\ & \quad + 5M \sqrt{X(\bar{x}(t), \bar{Z}(t), \bar{W}(t), \bar{\Lambda}^1(t), \bar{\Lambda}^2(t))} \end{aligned} \quad (27)$$

It can be easily verified that $X(\bar{x}(t), \bar{Z}(t), \bar{W}(t), \bar{\Lambda}^1(t), \bar{\Lambda}^2(t))$, $t \geq 0$, is bounded, so are $\bar{x}(t), \bar{Z}(t), \bar{W}(t), \bar{\Lambda}^1(t), \bar{\Lambda}^2(t)$ for all $t \geq 0$. As the result, the trajectory $(x, Z, W, \Lambda^1, \Lambda^2, \phi)$ is bounded.

ii) Let

$$\begin{aligned} \mathcal{R} & \subset \{ \phi \in \mathbb{R}_{(\tau m+1)nq} : x = P_{\Omega}[\bar{x}], x^* = P_{\Omega}[\bar{x}^*], \\ & \quad \min_{f_x \in \partial f(x), f_{x^*} \in \partial f(x^*)} (x - x^*)^T (f_x - f_{x^*}) = 0, \quad (28) \\ & \quad L_{mnq} \Lambda^1 = \mathbf{0}_{mnq}, L_{mnq} \Lambda^2 = \mathbf{0}_{mnq} \} \end{aligned}$$

Note that $(x - x^*)^T (f_x - f_{x^*}) > 0$ if $x \neq x^*$ since the Assumption 3.1. Hence, $\mathcal{R} \subset \{ \phi \in \mathbb{R}_{(\tau m+1)nq} : L_{mnq} \Lambda^1 = \mathbf{0}_{mnq}, L_{mnq} \Lambda^2 = \mathbf{0}_{mnq}, x = P_{\Omega}[\bar{x}] = x^* \}$. Let \mathcal{M} be the largest weakly invariant subset of \mathcal{R} . According to Lemma 2.3, $\phi \rightarrow \mathcal{M}$ as $t \rightarrow \infty$. Hence, $x(t) \rightarrow x^*$ as $t \rightarrow \infty$. Part (ii) is thus proved. \square

6 Simulation

In this section, we show a numerical example to validate our proposed distributed optimization algorithm. Consider the distributed robust optimization problem with four agents moving in a 2-D space with first-order dynamics as follows

$$F(x) = \sum_{i=1}^4 \|x_i - p_i\|_2^2 + |x|_1 \quad (29)$$

where $p_i = [i, -i]^T$, $\|\cdot\|_2$ denotes the l_2 norm, $\Omega_i = \{ \delta \in \mathbb{R}^2 : \|\delta - x_i(0)\|_2 \leq 30 \}$, $m = 2$, $\gamma_1 = \gamma_2 = 2$, $A_{i1} = 0.1 \cdot i \cdot I_2$, $\hat{A}_{i1} = 0.1 \cdot (5 - i) \cdot I_2$, $A_{i2} = \hat{A}_{i1}$, $\hat{A}_{i2} = A_{i1}$, and $b_1^1 = [-15, -5]^T$, $b_1^2 = [-10, -4]^T$, $b_1^3 = [0, -6]^T$, $b_1^4 = [4, 0]^T$, $b_2^1 = [-5, -1]^T$, $b_2^2 = [-4, -3]^T$, $b_2^3 = [0, -2]^T$, $b_2^4 = [1, -5]^T$, $b_1 = [-21, -15]^T$, $b_2 = [-8, -11]^T$. This problem can be transferred to its corresponding dual problem as the form of problem (5). The Laplacian of the undirected




Fig. 1: The trajectories of $x_i(t)$, $i \in \{1, 2, 3, 4\}$ with algorithm (8)




Fig. 2: The trajectories of $G_{j1}(x)$ and $G_{j2}(x)$, $j \in \{1, 2\}$ with algorithm (8)

graph \mathcal{G} is given by

$$L_4 = \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \quad (30)$$

The initial positions of the agents 1, 2, 3, and 4 are set as $x_1(0) = [-13, 12]^T$, $x_2(0) = [17, 15]^T$, $x_3(0) = [-10, -11]^T$ and $x_4(0) = [16, -14]^T$. We set the initial values for the Lagrangian multipliers λ_{ij}^1 , λ_{ij}^2 , μ_{ij} and auxiliary variables z_{ij} , w_{ij} , y_{ij}^1 , y_{ij}^2 as zeros for $i \in \{1, 2, 3, 4\}$, $j \in \{1, 2\}$. The optimal solution is $x_1^* = [-7.439, -10.408]^T$, $x_2^* = [-4.016, -6.409]^T$, $x_3^* = [-15.516, -17.612]^T$, $x_4^* = [-13.401, -19.965]^T$.

Fig.1 gives the trajectories of $x_i(t)$, $i \in \{1, 2, 3, 4\}$. It can be seen that the trajectory of x converges to the optimal solution. Let $G_{j1}(x) = \sum_{i=1}^4 H_{ij}^1$, $G_{j2} = \sum_{i=1}^4 H_{ij}^2$, $j \in \{1, 2\}$. Fig.2 shows the trajectory of $G_{j1}(x)$ and $G_{j2}(x)$, $j \in \{1, 2\}$, which proves that the constraint condition of problem (4) are satisfied.

7 Conclusion

In this paper, a distributed nonsmooth resource allocation problem with cardinality constrained uncertainty has been investigated. With the help of duality theory about convex optimization, a deterministic distributed robust resource allocation problem with linear optimization formulation has been derived under the framework of multi-agent system. A distributed projection-based algorithm has been proposed to deal with this problem. Based on stability theory and differential inclusions, the proposed algorithm has been proved to reach the optimal solution and satisfy the resource allocation condition simultaneously.

References

[1] R. Tibshirani, M. Saunders, S. Rosset, J. Zhu and K. Knight, "Sparsity and smoothness via the fused LASSO," *Journal of*

The Royal Statistical Society Series B: Statistical Methodology, 67(1): 91–108, 2005.

[2] C. Wu, H. Fang and X. Zeng, "Distributed object transport of mobile manipulators with optimal manipulable coordination," in *Proceedings of 37th Chinese Control Conference*, 2018: 7043–7048.

[3] S. Kai, H. Fang, C. Wu, X. Zeng and Q. Yang, "Distributed formation and motion control multiple mobile manipulator transportation: an energy optimization design," in *Proceedings of 37th Chinese Control Conference*, 2018: 1179–1184.

[4] A. Nedić and A. Ozdaglar, "Distributed subgradient methods for multi-agent optimization," *IEEE Transactions on Automatic Control*, 54(1): 48–61, 2009.

[5] T. H. Chang, A. Nedić and A. Scaglione, "Distributed constrained optimization by consensus-based primal-dual perturbation method," *IEEE Transactions on Automatic Control*, 59(6), 1524–1538, 2014.

[6] X. Zeng, P. Yi and Y. Hong, "Distributed algorithm for robust resource allocation with polyhedral uncertain allocation parameters," *Journal of Systems Science and Complexity*, 31(1): 103–119, 2018.

[7] S. Lee and M. M. Zavlanos, "Approximate projection methods for decentralized optimization with functional constraints," *IEEE Transactions on Automatic Control*, 63(10), 3248–3260, 2018.

[8] A. Cherukuri and J. Cortés, "Distributed generator coordination for initialization and anytime optimization in economic dispatch," *IEEE Transactions on Control of Network Systems*, 2(3): 226–237, 2015.

[9] P. Yi, Y. Hong and F. Liu, "Distributed gradient algorithm for constrained optimization with application to load sharing in power systems," *System and Control Letter*, 83: 45–52, 2015.

[10] S. Liang, X. Zeng and Y. Hong, "Distributed sub-optimal resource allocation over weight-balanced graph via singular perturbation," *Automatica*, 95: 222–228, 2018.

[11] A. Ben-Tal, L. El Ghaoui and A. Nemirovski, *Robust optimization*, Princeton: Princeton University Press, 2009.

[12] D. Bertsimas and M. Sim, "The price of robustness," *Operations Research*, 52(1): 35–53, 2004.

[13] Z. Li, R. Ding and C. A. Floudas, "A comparative theoretical and computational study on robust counterpart optimization: I. Robust linear optimization and robust mixed integer linear optimization," *Industrial and Engineering Chemistry Research*, 50(18): 10567–10603, 2011.

[14] H. Ghasvari, M. A. Raayatpanah and P. M. Pardalos, "A robust optimization approach for multicast network coding under uncertain link costs," *Optimization Letters*, 11(2): 429–444, 2017.

[15] Q. Liu and J. Wang, "A one-layer projection neural network for nonsmooth optimization subject to linear equalities and bound constraints," *IEEE Transactions on Neural Networks and Learning Systems*, 24(5): 812–824, 2013.

[16] F. H. Clarke, *Optimization and Nonsmooth Analysis*, New York: Wiley, 1983.

[17] J. Cortés, "Discontinuous dynamical systems," *IEEE Control Systems Magazine*, 44(11): 1995–2006, 1999.

[18] D. Bertsimas, D. B. Brown and C. Caramanis, "Theory and applications of robust optimization," *SIAM review*, 53(3): 464–501, 2011.

[19] A. Ruszczyński, *Nonlinear Optimization*. Princeton: Princeton University Press, 2006.